

Melanogaster minysporus sp. nov., a new sequestrate member of the Boletales from Mexico

Efrén Cázares¹, Gonzalo Guevara²,
Jesús García², James M. Trappe¹

¹ Oregon State University, Department of Forest Ecosystem and Society, Corvallis, Oregon 97331-5752, USA.
² Instituto Tecnológico de Cd. Victoria, Av. Portes Gil 1301 Pte. C.P. 87010, A.P. 175 Cd. Victoria, Tam., México

Melanogaster minysporus sp. nov. un nuevo miembro secuestrado de los Boletales de México

Resumen. Se describe una nueva especie *Melanogaster minysporus* recolectada en el norte de México, en el estado de Nuevo León. La especie se distingue por presentar un peridio amarillo brillante que se mancha de rojizo cuando se maltrata y por las esporas pequeñas.

Palabras clave: Paxillaceae, hipogeos, *Quercus*.

Abstract. *Melanogaster minysporus* is described as a new species from northern Mexico (Nuevo Leon). It is distinctive in the genus in having a bright yellow peridium that stains reddish where bruised in fresh and small spores.

Key words. Paxillaceae, hypogeous, *Quercus*.

Recibido 10 de octubre 2008; aceptado 5 de diciembre 2008.

Received 10 October 2008; accepted 5 December 2008.

Introduction

Carl Lumholtz, a Norwegian explorer, collected *Melanogaster variegatus* var. *mexicanus* Farlow nom. nud. (= *M. umbrinigleba* Trappe & Guzmán) while exploring the Chihuahuan mountains in northern Mexico (Lumholtz, 1902). This was the first record of a sequestrate fungus from Mexico. Cázares *et al.* 1992 described *Melanogaster variegatus* found in oak forests in northern Mexico. Here we describe another new *Melanogaster* species from northern Mexico. It is notable for its bright yellow peridium and small spores.

The methods of Castellano *et al.* (1989) were used for the morphological description. Hand-cut sections were mounted in 5% KOH and Melzer's reagent for examination by compound- and stereomicroscopy.

Results

Melanogaster minysporus Cázares, Guevara, García & Trappe, sp. nov.

Figures 1-3

Basidiomata 13-35 x 10-27 mm, *globosa* vel *subglobosabolbata*. *Peridium flavissimum*, *ubi contusum rubescens*, 160-240 µm *crassum*, *hyphis intertextis*. *Gleba solida*, *gelatinosacatilaginea*, *maturitate brunneola* vel *brunnea venis albis* vel

Autor para correspondencia: Efrén Cázares
mycoroots@comcast.net



Figures 1-3. *Melanogaster minysporus*. 1: Basidiomata (bar = 1 cm), 2: Basidiomata in cross section (bar = 1 cm), 3: Basidiospores (bar = 10 µm).

flavis marmorata. Sporae subfusiformes vel obovoideae, (5-) 5.5-6.5 x (3-) 4-5 µm, laeves, avellaneae vel brunnea.

Holotypus hic designatus: Mexico, Nuevo Leon, Santiago, El Barro, Guevara 907, ITCV.

Basidiomata hypogeous, 13-35 x 10-27 mm, globose to subglobose-lobed, depressed at the base, the surface smooth to finely fibrillose, bright yellow, staining reddish where bruised when fresh. Gleba solid, gelatinous-rubbery, pale brown in youth, at maturity dark brown with white to yellow, sterile veins. KOH reddish brown on fresh peridium and black on dried specimens. FeSO₄ black on dried specimens.

Peridium 160-240 µm thick. Epicutis of appressed, hyaline, thick-walled up to 1 µm thick, interwoven hyphae upto 4 µm in diam with clamp connections, orange-brown to

yellow-brown in KOH and Melzer's reagent. Subcutis of interwoven hyphae, thin- to thick walled, hyphae hyaline in KOH and yellowish in Melzer's reagent. Spores (5-) 5.5-6.5 x (3-) 4-5 µm, subfusoid to obovate, smooth, thick-walled, up to 1 µm thick, the sterigmal attachment distinct, in KOH and Melzer's reagent yellow brown to brown singly, dark brown to black in mass.

Etymology: Greek, *miny-* (small) and *spora* (spores).

Habitat and season: Hypogeous to subhypogeous, single to gregarious, in mixed *Pinus-Quercus* forests, under *Quercus canbyi* and *Q. rhizophylla*. August through September.

Collections examined: HOLOTYPE: Mexico, Nuevo Leon, Santiago, El Barro, camino al paraje del oso, 6

August, 2007, Guevara 907 (ITCV, isotype OSC). PARATYPES (all ITCV): Nuevo Leon, Sierra Madre Oriental, Mpio. Santiago, Cañon de Puerto Genovevo, 11 September 1983, García 3802; 7 September 1983, García 3803; 7 August 1983, García 3804; 7 August 1983, García 3805; 29 September 1983, García 3814; Mpio. Guadalupe, La Pastora, 20 June 1984, G. Valencia s.n.

Discussion. This species is distinguished by its bright yellow peridium that stains reddish where bruised when fresh, KOH and FeSO₄ black on dried peridium, and small spores. *Melanogaster ovoidisporus* Y. Wang, described from China (Wang *et al.*, 1995) is a similar species with small spores (5-) 5.5-6.5 (-7) x (3.5-) 4-5 (-5.8) µm, but its peridium is brown. They also described *M. ovoidisporus* var. *angustatosporus* K. Tao, Chang & Liu with narrower spores 5.5-6.5 (-7) x 3-4 µm.

Alpova microspora (Velen.) Trappe, has a yellow peridium and a rubbery-gelatinous gleba, but its spores are fusiform to oblong, (4-) 4.5-6 (-7) x 2-3 (-4) µm and its gleba

stains red when exposed; this species was described from central Europe (Trappe, 1975).

Acknowledgments

We thank the Dirección General de Educación Superior Tecnológica for financial support.

References

- Castellano, M.A., J.M. Trappe, Z. Maser, C. Maser, 1989. Key to spores of the genera of hypogeous fungi of north temperate forests with special reference to animal mycophagy. Mad River Press, Eureka, California.
- Cázares, E., J. García, J. Castillo, J. M. Trappe, 1992. Hypogeous fungi from northern Mexico. Mycologia 84: 341-359.
- Lumholtz, C. 1902. Unknown Mexico. Vol 1. Charles Scribner's Sons, New York.
- Trappe, J. M. 1975. A revision of the genus *Alpova* with notes on *Rhizopogon* and the Melanogastraceae. Nova Hedwigia Beiheft 51: 279-310.
- Wang, Y., Chang, M., Tao, K., Liu, B. 1995. New species and new varieties of the genus *Melanogaster* from Ching. Journal of Shanxi University (Nat. Sci. Ed.) 18: 449-453.